



1/12



5 Figure 1: XRD pattern of composite following co-precipitation (Cu-K α radiation).

10



15 Figure 2: SEM micrograph of triple co-precipitate.

2/12




Figure 3: TEM micrograph of triple co-precipitate




Figure 4: XRD pattern of composite following dehydrothermal treatment at 105°C and 50mTorr for 48 hours, indicating that the brushite phase has converted to its dehydrated form monetite.

3/12




Figure 5: set of combinations of ionic concentration and calcium nitrate: calcium hydroxide ratio for maintaining pH = 4.0.

5







Figure 6: Identification of conditions for pH 4.0 synthesis of a triple coprecipitate slurry containing a 1:1 mass ratio of calcium phosphate to collagen plus GAG.

10

4/12




5 Figure 7: x-ray diffraction pattern of collagen/GAG/brushite triple coprecipitate following removal of unbound water (Cu- K_{α} radiation).




10 Figure 8: Secondary (SE) and backscattered electron (BSE) images of surface of triple coprecipitate with CaP: collagen + GAG = 1:1.

5/12



5 Figure 9: x-ray diffraction pattern of collagen/GAG/brushite triple coprecipitate following EDAC crosslinking (Cu- K_{α} radiation).



10 Figure 10: x-ray diffraction pattern of EDAC-crosslinked collagen/GAG/CaP triple coprecipitate following conversion at 37°C to octacalcium phosphate (OCP) over 72 hours at pH 6.67, to form a
15 collagen/GAG/OCP biocomposite (Cu- K_{α} radiation).

6/12




Figure 11: set of combinations of ionic concentration and calcium nitrate: calcium hydroxide ratio for maintaining pH = 4.5.

5






Figure 12: Identification of conditions for pH 4.5 synthesis of a triple coprecipitate slurry containing a 3:1 mass ratio of calcium phosphate to collagen plus GAG.

10

7/12



5 Figure 13: x-ray diffraction pattern of collagen/GAG/brushite triple coprecipitate following removal of unbound water (Cu- K_{α} radiation).




10 Figure 14: x-ray diffraction pattern of collagen/GAG/brushite triple coprecipitate following EDAC crosslinking (Cu- K_{α} radiation).


15

20

8/12



5 Figure 15: x-ray diffraction pattern of EDAC-crosslinked collagen/GAG/CaP triple coprecipitate following conversion at 37°C to apatite over 72 hours at pH 8.50, to form a collagen/GAG/apatite biocomposite (Cu- K_{α} radiation).




10 Figure 16: X-ray diffraction pattern of EDAC-crosslinked collagen/GAG/Ap triple coprecipitates after secondary crosslinking via gamma irradiation.

9/12

5

10



30

Figure 17


35

40

45

10/12

5



50 μm

10

Figure 18

15

11/12

5




Figure 19

30

35

40

45

12/12

5

10

15

20

25

30

35

40




Figure 20

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ BLACK BORDERS

☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

☒ FADED TEXT OR DRAWING

☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING

☐ SKEWED/SLANTED IMAGES

☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS

☐ GRAY SCALE DOCUMENTS

☐ LINES OR MARKS ON ORIGINAL DOCUMENT

☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.